

Bovine respiratory syncytial virus *in-situ* hybridization from sheep lungs at different times postinfection

Hibridación *in situ* del virus respiratorio sincytial bovino en pulmón de cordero a diferentes tiempos postinfección

E. REDONDO¹, D.V.M., Ph.D.; L. GOMEZ¹, D.V.M., Ph.D.; C. L. KELLING², D.V.M., Ph.D.;
A. GAZQUEZ¹, D.V.M., Ph.D.; A.J. MASOT¹ D.V.M., Ph.D.

¹Histology and pathology, Faculty of Veterinary Medicine, University of Extremadura, 10071 Cáceres, Spain.

²Department of Veterinary and Biomedical Sciences, University of Nebraska-Lincoln, USA.

e-mail: eloy@unex.es

RESUMEN

Se estudió, mediante la técnica de hibridación *in situ*, la distribución del ARN viral del virus respiratorio Sincicial Bovino (VRSB) en pulmón de corderos infectados en forma experimental, a diferentes tiempos postinoculación. La sonda usada para la hibridación *in situ* se preparó mediante transcripción reversa del ARN del VRSB, seguida de la amplificación mediante PCR de cADN. 25 corderos de raza Merino de ambos sexos y de un peso vivo de 55 (+/- 10) Kg, fueron inoculados por vía intratraqueal con 40 ml de solución salina que contenía 1.26×10^6 DIM₅₀ por ml (cepa viral NMK7). Los corderos se sacrificaron los días 1, 3, 7, 11 y 15 postinoculación. Las células epiteliales bronquiales y bronquiales resultaron positivas a los ácidos nucleicos de VRSB los días 1, 3, 7 y 11 postinoculación. A su vez, el epitelio alveolar contenía células positivas los días 1, 3, y 7 postinoculación. Se detectaron células que contenían el ARN viral en las luces bronquiales y bronquiales del día 1 al 11 postinoculación, y en el exudado alveolar en los días 3 y 7 postinoculación. Se identificaron señales positivas de hibridación desde el día 3 al 11 postinoculación, tanto en las células intersticiales mononucleares como en el tejido linfoido asociado a los bronquios. Las señales de hibridación más intensas se detectaron a los días 3 y 7 postinoculación, lo que coincidió con las lesiones histopatológicas de mayor consideración.

Palabras claves: Virus respiratorio sincicial bovino, hibridación, oveja, pulmones.

Key words: bovine respiratory syncytial virus, *in situ* hybridization, sheep, lungs.

INTRODUCTION

Respiratory syncytial virus (RSV) is a respiratory tract pathogen of both humans and cattle. Human RSV is the most important cause of viral lower respiratory tract disease in infants and young children (Bruhn and Yeager, 1977). Bovine and human RSV strains (enveloped RNA virus) are pneumoviruses of the Paramyxoviridae family and are currently considered to be

antigenically related (Thomas *et al.*, 1984). Although greater susceptibility has been reported in cattle (Castleman *et al.*, 1985; Bryson *et al.*, 1988; 1991a,b; Kliman *et al.*, 1989; Oberst *et al.*, 1993), BRSV may also affect other domestic species, particularly small ruminants (Al-Darraj *et al.*, 1982; Bryson *et al.*, 1988; Masot *et al.*, 1993 a,b; 1995; 1996; Redondo *et al.*, 1994; Belknap *et al.*, 1995).

The RSV fusion protein causes fusion of viral and host cellular membranes and cell to cell fusion (Lerch *et al.*, 1991). This protein is synthesized as F₀ precursor, which is

proteolytically cleaved into two disulfide-linked polypeptide subunits, F₁ (48KDa) and F₂ (20KDa), that form the active fusion protein (Walsh *et al.*, 1985; Walravens *et al.*, 1990; Lerch *et al.*, 1991; Paton *et al.*, 1992; Pasty and Samal, 1993; Matheise *et al.*, 1995). The BRSV F mRNA is 1899 nucleotides in length (Lerch *et al.*, 1991). The nucleotide sequence encoding the F protein is highly conserved (>95%) among BRSV strains (Pasty and Samal, 1993). This identity is lower between F mRNAs of RSV of bovine and human origin (71.5%) (Lerch *et al.*, 1991).

A digoxigenin-labeled DNA probe corresponding to the part of the genome that codes for bp 736-1522 of the BRSV protein gene was developed for localization of virus in lung sections.

The aim of the present study is to analyze the pathology of sheep experimentally infected with BRSV, and to visualize specific hybridization signals for this virus in lung samples.

MATERIALS AND METHODS

Animals and BRSV inoculum. Twenty-five gnotobiotic merino sheep of both sexes with a live weight of 55 ± 5 Kg, each received a intratracheal inoculation of 40 ml of the NMK7 strain of BRSV (supplied by Dr. Gómez-Tejedor, Instituto Nacional de Investigaciones Agrarias, Madrid, Spain) diluted to a concentration of 1.26 x 10⁶ TCID₅₀/ml. The origin and characterization of the NMK7 isolate of BRSV has been previously reported by Belknap *et al.*, (1995). The sheep were housed in an isolation barn, and were euthanized 1,3,7,11 and 15 postinoculation days (PID).

Virus inoculum was prepared using primary bovine fetal kidney (BFK) cell cultures infected with a BRSV stock strain NMK7, to which minimum Essential medium (Sigma, Madrid, Spain) with 15% bovine fetal serum (Sigma, Madrid, Spain) was added; the final cultures were incubated at 37°C in a CO₂ atmosphere. Ten animals used as control were inoculated with an identical volume of un-infected BFK cell culture. They were kept in another isolation box and were slaughtered 1,3,7,11 and 15 PID.

Aliquots of virus were cultured and were determined to be free of aerobic bacteria and mycoplasma by standard techniques.

Samples of lung were collected for microbiological studies and virus isolation. Formalin fixed sections were used for histopathological techniques and in-situ hybridization assay.

Histopathology. Sections, 3 µm thick, were cut and stained with haematoxylin and eosin for routine morphological studies.

Immunoperoxidase staining procedure for detection of BRSV antigens. An Avidin-Biotin-Peroxidase complex was carried out on deparaffined and trypsinized lung samples. Sections were blocked in diluted (1:50) normal swine serum (Dakopatt, Spain) for 15 minutes to reduce background and were incubated in diluted (1:1000) rabbit anti-RSV (Dakopatt, Spain) for 3 hours at 20°C. Diluted (1:500) biotinylated swine anti-rabbit IgG (Dakopatt, Spain) was placed on the sections for 30 minutes, followed by 1 hour incubation in diluted (1:50) ABPC reagent (Dakopatt, Spain). Sections were incubated in diaminobenzidine solution for 5 minutes and were counterstained lightly with Mayer's haematoxylin. The positive controls included BRSV-infected BFK cell cultures (the cytopathic effect was evident in the formation of numerous syncytia) and noninfected BFK cell cultures. Test control sections were also stained using nonimmune rabbit serum as first layer.

Microbiological investigations. Lung samples tissue were homogenized in phosphate-buffered saline and a serial 10-fold dilutions were made up to 10⁴ and 0.1 ml of each dilution spread on to half a sheep's blood agar plate. Duplicate plates were used, one incubated aerobically and the other anaerobically under H₂ containing 10% CO₂. Colonies were identified using the methods described by Cowan (1974). Bronchial swabs and lung samples were also cultured for mycoplasma in modified Hayflick's broth, and colonies were identified on agar plates.

Nasal swab specimens and lung samples were used to detect BRSV by virus isolation in BFK

cells, following the technique described previously by Castleman *et al.*, 1985. Isolates were confirmed as BRSV by indirect immunofluorescence (Masot *et al.*, 1993a).

RNA extraction. BT cells (75 cm² flasks) were infected with BRSV strain NMK7 (Lerch *et al.*, 1991; Oberst *et al.*, 1993); provided by Dr. Kelling, Department of Veterinary and Biomedical Sciences, University of Nebraska, Lincoln, USA. This is a BRSV strain plaque purified more than three times that was originally obtained from cattle and is bovine viral diarrhoea (BVD) virus free (Lehmkuhl *et al.*, 1979). Cell cultures were incubated at 33 °C in high glucose Dulbecco modified Eagle medium (DMEM) supplemented with 2% horse serum. Viruses were allowed to replicate and form syncytia (cytopathic effect), at which point the RNA was extracted from infected cells. RNA was isolated with the Trizol LS Reagent (Gibco, NY, USA) following the manufacturer's instructions. RNA concentration of the extracts was estimated by measuring the A₂₆₀ with a Beckman DU-64 Spectrophotometer.

RT-PCR. An amplified product of 786 bp length encoding for bp 736-1522 of the BRSV protein F gene, was obtained by nested RT-PCR. Primer 1555 (5' TTTCTACATCTACTG 3') was used to obtain cDNA and primers 736 (5' TGGTATTACCACCCCCTCAG 3') and 1522 (5' TCAGATCGACGTATGAAAGCC 3') were used for PCR. The PCR reaction was performed in 35 cycles of 1 min at 95 °C, 1 min at 55 °C and 1 min at 72 °C.

Cloning and labeling probe. The amplified product was cloned in the pCR 2.1 plasmid vector (Invitrogen, CA, USA). The correct sequence was confirmed by sequencing.

The probe was prepared by removing the DNA insert by digestion with restriction enzyme *EcoRI*. The DNA was submitted to gel electrophoresis and the fragment extracted using Qiaex II (Quiagen, Hilden, Germany). DNA fragment was digoxigenin-labeled by random priming (Boehringer Mannheim Biochemicals,

Mannheim, Germany). The specificity of the probe was checked by dot blot and southern blot onto nylon membranes.

Preparation of samples for ISH. BRSV-infected and uninfected cell cultures were used for optimization of ISH. BT cells were infected and grown as has been described before. The cultures that showed cytopathic effect were trypsinized and washed twice with DMEM. The first time with the DMEM containing 5% equine serum and the second time containing as serum. After counting, the harvested cells were applied to ProbeOn Plus microscope slides (Fisher Scientific, Pittsburgh, PA, USA) by centrifugation at 1500 rpm for 3 min (cytocentrifuge Shandon Cytospin 2) at a density of 2x10⁵ cells per spot. Slides were fixed in 4% paraformaldehyde and dehydrated through an ethanol series.

Lung sections, 3 µm thick, were deparaffinized with xylene and rehydrated through decreasing concentrations of ethanol. All the samples (slides containing cells or sections) were equilibrate in phosphate-buffered saline (PBS). Permeabilization to allow the penetration of the probe was carried out by treatment with 0.2N HCl for 20 min at room temperature and digestion with 5 µg/ml of proteinase K (Promega, Madison, WI, USA) for cells and 30 µg/ml for tissues in PBS for 20 min at 37°C. Some slides were treated with 50 µg/ml of RNase (Boehringer Mannheim Biochemicals) in PBS for 60 min at 37°C in wet chamber, to be used as specificity control. Next, slides were refixed in 5% paraformaldehyde and followed by two rinses in PBS. Slides were acetylated in 100 mM triethanolamine with 0.25% of acetic anhydride (to neutralize positive charges and thus reduce nonspecific electrostatic binding of the probe) for 10 min. Slides were then washed twice in 2x SSC (1x SSC is 15 mM of sodium citrate plus 0.15 M NaCl, pH 7.0).

Prehybridization. Slides were prehybridized at 70°C for 60 min in a humidified chamber by adding 200 µl of hybridization solution containing 4x SSC, 2mM EDTA, 10% dextran

sulfate, 50% deionized formamide, 500 µg of salmon sperm DNA and 1x Denhardt solution (0.02% polyvinylpyrrolidone, 0.02% bovine serum albumin and 0.02% Ficoll 400) per ml, without cDNA probe.

Hybridization. The ISH conditions were as described elsewhere (Sur *et al.*, 1996).

Enzyme-Linked Immunoabsorbent Assay (ELISA). Positive control antigen was prepared from bovine foetal kidney cell cultures infected with BRSV strain NMK7. Negative control antigen was determined by block titration; those which yielded maximum optical density (OD) values for positive sera without causing an increased OD in negative sera were considered satisfactory.

Polystyrene plates were coated with 100 µl of the antigen diluted to a concentration of 5µg/ml in sodium carbonate buffer (0.1 M, pH 9.6) with 0.02% NaN₃ for 3 h at 37 °C. After coating and between all subsequent steps, the plates were washed 4 times in phosphate-buffered saline containing 0.05% Tween 20 (PBSTw). As a blocking agent, 5% bovine serum albumin (BSA) in washing buffer was added (200ml) and incubated for 30 min at 37 °C. After washing, serum samples diluted to 1/1000 in PBSTw (100 µl) were added (200ml) and incubated for 30 min at 37 °C. After subsequent washing, 100 µl of horseradish peroxidase conjugate mouse anti-sheep IgG (diluted 1/2000 in washing buffer) were added. The reaction was developed at room temperature by the addition of 100 µl of 0.04% ophenylenediamine and 0.004% H₂O₂ in citrate-phosphate buffer (pH 5), stopped after 1h by the addition of 50 µl of 3N H₂SO₄ and read on a spectrophotometer at 490 nm.

The O.D. of BRSV = O.D of BRSV Ag – O.D. Control Ag.

Light Nephelometry. A biological fluid protein counter was used to determine immunoglobulin levels. This system measures the intensity of light and the dispersion of the light beam as it passes through particles in cell fluid suspension. Particles were formed by

immunoprecipitation of diluted (1:100) experiment sera with diluted (1:50) monoclonal sheep anti-IgA, anti-IgG and anti-IgM antibodies. Immunoglobulin levels were expressed in mg/dl.

RESULTS

BRSV inoculum. Viral inoculum was free of aerobic bacteria, mycoplasma and BVD virus.

Histopathology. The histopathological study has been previously reported by us on studies reported here (Masot *et al.*, 1995, 1996). Catarrhal bronchiolitis observed on 1 PID was associated with granulocyte infiltration of the bronchiolar lumen. The interalveolar septa were thickened, with pronounced interstitial edema and moderate cell reaction.

Alveolar exudate in animals slaughtered 3 and 7 PID consisted of neutrophils, lymphocytes and multinucleate giant cells which formed the syncytia. There was considerable thickening of the interalveolar septa, due to the presence of edema and granulocyte and monocyte infiltration. Bronchiolitis was accompanied by epithelial cell necrosis. Hyperplasia of bronchiolar epithelial cells was conspicuous and early stages of reepithelization were apparent. Exudate in the bronchiolar lumina consisted of desquamated cells, necrotic debris and syncytial cells.

Severe bronchial, bronchiolar and alveolar damage was visible by 11 PID. Lung parenchyma had clear focal areas of consolidation due to bronchiolitis obliterans and the alveolar collapse caused by infiltration of macrophages, lymphocytes and syncytial cells into lumina.

Animals slaughtered at 15 PID presented a marked interstitial inflammatory reaction, with considerable septal thickening. Intense bronchiolar and/or alveolar hyperplasia was also observed. Lung consolidation was less marked than in animals slaughtered earlier.

IHC signals. The IHC study has been previously published by us on studies reported here (Masot *et al.*, 1993b, 1996). BRSV antigen was detected in bronchial and bronchiolar

epithelial cells, in bronchial mucous cells and in alveolar epithelial cells at 3 and 7 PID. Intense staining was also observed in alveolar macrophages, interstitial mononuclear cells and syncytia from 3 to 11 PID. Antigen was commonly detected in exudate within bronchiolar, bronchial and alveolar lumina. Specific staining was absent in the negative control (table 1).

Microbiological findings. Twenty three of the sheep inoculated with BRSV and nine of the control animals, were bacteriologically negative at the conclusion of the experiment. Contaminative species of bacteria were *E. coli* and *Bacillus spp.* Mycoplasma was not isolated from the upper or lower respiratory tract of any sheep.

ISH signals. In all PID the specificity of the signal reaction was confirmed by the rigorous observance of specific controls (figure 1, 2). The BRSV nucleic acid signal was specific because this was completely absent from non infected cells (figures 1, 2). There was no detectable ISH signal of BRSV RNA in the lungs of the control sheep (table 1).

Hybridization signals specific for BRSV RNA were detected in bronchial and bronchiolar epithelial cells at 1,3,7 and 11 PID (figure 3 and 4) but they were not observed at 15 PID.

BRSV-positive cells was commonly detected in exudate within bronchial and bronchiolar lumina from 1 to 11 PID (figure 5).

The positive IHS signal was observed in alveolar epithelial cells at 1, 3 and 7 PID (figures 6, 7, 8). In several cases the signal of RNA-virus also coincided with a cell morphology and location that would suggest that the cell could be classified as a type II pneumocyte. The RNA genome was visible in: free cells in alveolar space (figure 9) and in syncytial alveolar cells (figure 10) at 3 and 7 PID.

Hybridization signal specific for BRSV RNA was observed in the peribronchiolar area (figures 11,12), in bronchus associated lymphoid tissue (figure 12) and in mononuclear cells of interalveolar septum from 3 to 11 PID.

ELISA. Serum IgG BRSV antibody titres are summarised in figure 13. A progressive increase in titres was recorded until 7 PID; giving way, to an decrease from 11 PID onwards. The highest antibody response was observed in PID 3 and 7.

Light Nephelometry. Serum IgA, IgM and IgG levels are shown in table 2. The maximum concentrations of IgA and IgG were detected in 11 and 15 PID. From 1 until 7 PID, similar values of IgA and IgG were detected in sera of control and experimental animals. IgM levels remained above those recorded for preinfected animals through the experiment.

Viral isolation

Results of BRSV isolation attempts from lung homogenates or nasal secretions are presented in table 2.

DISCUSSION

The positive IHS signal in bronchi and bronchioles is indicative of the direct pathogenic action of BRSV at these levels (Viuff *et al.*, 1996) inducing necrosis and desquamation of bronchial and bronchiolar epithelia. Necrosis of bronchial and/or bronchiolar epithelia has been described elsewhere as a limitation of pulmonary antibacterial defenses with impaired mucociliary clearance, and as a predisposition to secondary bacterial infection (Pirie *et al.*, 1981; Al-Darraj *et al.*, 1982; Castleman *et al.*, 1985; Bryson *et al.*, 1988; Redondo *et al.*, 1994; Masot *et al.* 1993 a,b; 1995; 1996; Viuff *et al.*, 1996).

Cells obstructing bronchial and bronchiolar lumina tend to form syncytia (Bryson *et al.*, 1988; 1991a,b; Ciszewski *et al.*, 1991; Masot *et al.*, 1993a,b; 1995; 1996; Belknap *et al.*, 1995) which are usually located in bronchioles and rarely observed in bronchi. This finding has previously been reported in studies of immunohistochemistry (Masot *et al.*, 1993a,b) or *in situ* hybridization (Viuff *et al.*, 1996). The possible involvement of type I and II pneumocytes in BRSV replication in sheep lung has been detected by IHC (Masot *et al.*, 1993a,b, 1995, 1996). Extensive hypertrophy and

TABLE 1. IHC and ISH signal in the lung of experimentally infected sheep by BRSV.
IHC and ISH signal in the lung of experimentally infected sheep by BRSV.

		ISH								IHC							
		BEC	BE	B _r EC	B _r E	AEC	SYN	BALT	MC	BEC	BE	B _r EC	B _r E	AEC	SYN	BALT	MC
	Nº 26	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Nº 27	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Nº 28	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Nº 29	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Nº 30	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Nº 31	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Nº 32	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Nº 33	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Nº 34	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Nº 35	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Nº 1	+	+	+	+	+	+	+	+	+	+	+	+	+	-	-	-
	Nº 2	+	+	+	+	+	+	+	+	+	+	+	+	+	-	-	-
	Nº 3	+	+	+	+	+	+	+	+	+	+	+	+	+	-	-	-
	Nº 4	+	+	+	+	+	+	+	+	+	+	+	+	+	-	-	-
	Nº 5	+	+	+	+	+	+	+	+	+	+	+	+	+	-	-	-
	Nº 6	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	Nº 7	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	Nº 8	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	Nº 9	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	Nº 0	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	Nº 11	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	Nº 12	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	Nº 13	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	Nº 14	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	Nº 15	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	Nº 16	-	-	-	-	-	-	+	+	+	+	+	+	-	-	+	+
	Nº 17	-	-	-	-	-	-	+	+	+	+	+	+	-	-	+	+
	Nº 18	-	-	-	-	-	-	+	+	+	+	+	+	-	-	+	+
	Nº 19	-	-	-	-	-	-	+	+	+	+	+	+	-	-	+	+
	Nº 20	-	-	-	-	-	-	+	+	+	+	+	+	-	-	+	+
	Nº 21	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Nº 22	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Nº 23	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Nº 24	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Nº 25	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

+, Lamb positive by IHC or ISH signal. - Lamb negative by IHC or ISH signal.

BEC = bronchial epithelial cells; BE = bronchial exudation; B_rEC = bronchiolar epithelial cells; B_rE = bronchiolar exudation. AEC = Alveolar epithelial cells; SYNC = syncytial alveolar cells; BALT = bronchus associated lymphoid tissue; MO = Mononuclear cells.




FIGURE 1. Bovine Respiratory Syncytial Virus (BRSV) infected BT cells hybridized with a DIG-labelled cDNA probe. A positive ISH signal in the cytoplasm of cells is shown. Methyl green counterstain. Bar = 160 μ m.

FIGURE 2. *In situ* hybridization of mock-infected BT cells with the same probe. Methyl green counterstain. Bar = 160 μ m.

FIGURE 3. Lung of sheep BRSV-infected at 7 days p.i. ISH positive signal in bronchiolar epithelial cells. ISH counterstained with methyl green. Bar = 80 μ m.

FIGURE 4. Lung; sheep inoculated with BRSV and slaughtered at 11 PID. ISH positive signal in bronchiolar epithelial cells. ISH counterstained with methyl green. Bar = 80 μ m.

FIGURE 5. Lung; sheep inoculated intratracheally 7 days previously with BRSV. BRSV RNA infected cells located in bronchiolar exudations. ISH counterstained with methyl green. Bar = 40 μ m.

FIGURE 6. Lung; sheep inoculated intratracheally 1 days previously with BRSV. BRSV nucleic acid detected in alveolar epithelial cells. ISH counterstained with methyl green. Bar = 160 μ m.




FIGURE 7. Lung; sheep inoculated intratracheally 3 days previously with BRSV. ISH positive signal observed in alveolar epithelial cells. These cells that by morphology and localization could be classified as type II pneumocytes. ISH counterstained with methyl green. Bar = 80 μ m.

FIGURE 8. Lung; sheep inoculated intratracheally 7 days previously with BRSV. BRSV infected cells located in alveolar epithelium. ISH counterstained with methyl green. Bar = 80 μ m.

FIGURE 9. BRSV infected lung tissue taken from an infected sheep at 3 PID. Large numbers of positive cells free in alveolar space. ISH counterstained with methyl green. Bar = 80 μ m.

FIGURE 10. BRSV infected lung tissue taken from an infected sheep at 7 PID. Syncytia containing BRSV RNA positive cells. ISH counterstained with methyl green. Bar = 80 μ m.

FIGURE 11. Lung; sheep inoculated with BRSV and slaughtered at 3 PID. Positive signal in the peribronchiolar area. ISH counterstained with methyl green. Bar = 80 μ m.

FIGURE 12. Positive signal in the peribronchiolar area of the lung of sheep at 3 days postinoculation with BRSV. ISH counterstained with methyl green. Bar = 80 μ m.




FIGURE 13. Sheep experimentally infected with BRSV. Serum IgG BRSV antibody titres detected by ELISA. PID 1 (sheep number 1-5); PID 3 (sheep number 6-10); PID 7 (sheep number 11-15); PID 11 (sheep number 16-20); PID 15 (sheep number 21-25); Control: PID -10 (sheep number 26-30); PID -0 (sheep number 31-35).

hyperplasia of type II pneumocytes are also known to occur during acute BRSV infection in the lung (Masot *et al.*, 1993a, 1995, 1996). In the present study, results with IHS bears out the role of type II pneumocytes in BRSV replication, since the lung section showed cells exhibiting positive IHS signals and which were morphologically and topographically consistent with type II pneumocytes. In this connection, Viuff *et al.*, (1996), in natural infected calves, suggest that the replication of BRSV in the alveolar type II cells, may lead to a change in the amount and quality of surfactant. The deficiency of surfactant caused by BRSV replication in the type II pneumocytes, may be a determining factor in collapse of alveoli. The alveolar collapse was visible in lungs of experimentally infected sheep described in the present study. In order to confirm that type II pneumocytes represent a specific BRSV replication site, TEM should be performed on the same sections as those used for IHS.

In this experiment, a positive IHS signal for BRSV in alveolar macrophages could not be established (Viuff *et al.*, 1996). These results agree with previous reports (Schrijver *et al.*, 1994) suggesting that *in vitro* bovine alveolar macrophages (BAMs) exhibit a high intrinsic resistance for infection with BRSV and that bovine alveolar macrophages do not appear to be important for replication of BRSV. However, large numbers of BAMs may harbour virus antigen, even for several days, that may influence the function of these cells (Schrijver *et al.*, 1994). However, replication of human respiratory syncytial virus in alveolar macrophages has been shown (Cirino *et al.*, 1993).

A large number of alveolar syncytia containing BRSV-RNA were detected. The location of these syncytia coincided with the positive hybridization signals reported by Viuff *et al.*, (1996), and with the site of BRSV antigen detected elsewhere by IHC (Bryson *et al.*, 1988a; Masot *et al.*, 1993a,b, 1995, 1996). However, the reason for syncytia

TABLE 2. BRSV isolation from nasal secretions and lung homogenates. Means of serum immunoglobullin levels (g/dl). BRSV isolation from nasal secretions and lung homogenates. Means of serum immunoglobullin levels (g/dl).

BRSV ISOLATION		IgG (g/dl).	IgA (g/dl).	IgM (g/dl).
Lung	Nasal secretions			
-	-	2180	33	214
-	-	2183	35	216
-	-	2186	36	217
-	-	2190	32	220
-	-	2193	34	218
-	-	2188	35	221
-	-	2187	33	222
-	-	2183	34	221
-	-	2189	33	218
-	-	2190	32	219
-	-	2215	34	215
+	-	2220	33	222
+	+	2218	33	218
+	+	2216	34	221
+	+	2221	32	219
+	-	1967	36	228
+	+	1951	33	208
+	+	1957	37	211
+	+	1964	34	214
+	+	1967	32	217
+	+	1975	30	219
+	-	1980	33	224
+	+	1983	35	222
+	+	1986	36	226
+	+	1978	31	221
-	-	3245	46	212
-	-	3280	49	209
+	+	3275	51	211
-	+	3282	52	214
-	-	3269	50	204
-	-	3345	54	218
-	-	3359	51	222
+	+	3367	56	209
-	-	3298	59	216
-	-	3396	61	221

+, Lamb positive by viral isolation. -, Lamb negative by viral isolation.

formation remains an enigma. Syncytia are probably a result of fusion of cells and a direct spread of virus from cell to cell (Viuff *et al.*, 1996).

In the present study, positive ISH signals were observed in mononuclear cells in the interstitial (interalveolar and peribronchial) space and in BALT of damaged areas from 3 to 11 dpi. The distribution and topography of specific hybridization signals in the interstitium was very similar to the location of BRSV antigen detected by IHC (Bryson *et al.*, 1988a; Masot *et al.*, 1993c, 1995, 1996). However, the site of BRSV-RNA positive cells in BALT was a differentiating feature, since similar findings have not been reported elsewhere (Viuff *et al.*, 1996).

Interstitial cells expressing BRSV-RNA may carry virus particles to other organs, as described elsewhere (Lerch *et al.*, 1991).

The results obtained showed that more intense positive hybridization signals were detected at 3 and 7 dpi. It was precisely on these days that lesions were most intense (Masot *et al.*, 1993a,b, 1995, 1996), coinciding with maximum levels of serum IgG anti-BRSV. (Masot *et al.*, 1993a). However, this relationship is not clearly defined, since previous reports (Ciszewski *et al.*, 1991), state that levels of anti-BRSV serum antibodies remained relatively constant from 4 to 21 dpi with BRSV.

High values for IgG (PID 11 and 15) were associated with a marked decrease of BRSV antibody titres on days 11 and 15 PI, (Korbecki and Maksymowicz, 1977). Other authors report high neutralising antibody levels in calves from 5 PID onward, although maximal levels were detected at 5 weeks PI (Elazhary *et al.*, 1981; Castleman *et al.*, 1985). However, other studies have found low serum IgG values in experimentally-infected calves, although this response was recorded only when the animals were exposed to the virus for second time (Mohanty *et al.*, 1975). A marked increase in serum IgA levels were coincident with the decrease of virus antibodies titres over the same period (Korbecki and Maksymowicz). Other studies, however, report an absence of IgA in the serum of calves experimentally infected with human and bovine RSV (Thomas *et al.*, 1984).

In our experience, serum IgM levels did not increase. This finding conflicts with the increase in serum IgM levels described in children with RSV infection (Bruhn and Yeager, 1977; Korbecki and Maksymowicz, 1977).

In previously published experiments using parallel sections for IHC and IHS (Viuff *et al.*, 1996), authors were able to show, through IHC, that sites containing viral antigen corresponded to virus replication sites.

It is a well-known fact that immunodetection requires a large number of molecules in order to obtain positive reactions. In prolonged post-infection periods, the number of positive cells detected by immunohistochemistry (IHC) decreased; it was also sometimes difficult to differentiate between positive and negative signals, especially with high levels of background noise. IHC therefore has more limitations than IHS; with impaired antigen production in infected cells, viral RNA in cells persists (Schrijver *et al.*, 1994).

Further studies of comparison of consecutive serial sections are necessary to establish cell to cell correlation between the signal of both techniques.

SUMMARY

We studied the distribution of bovine respiratory syncytial virus (BRSV) RNA in lungs of experimentally infected sheep by *in situ*-hybridization at different times postinfection. The probe used for *in-situ* hybridization was prepared by reverse transcription of BRSV RNA, followed by PCR amplification of the cDNA. Twenty five Merino sheeps of both sexes with a live weight of 55 ± 10 Kg, received a intratracheal inoculation of 40 ml saline solution containing 1.26×10^6 TCID₅₀ BRSV (strain NMK7) per ml. Sheep were slaughtered 1,3,7,11 and 15 postinoculation days (PID). Bronchial and bronchiolar epithelial cells were positive for BRSV nucleic acid by ISH at 1, 3, 7 and 11 PID. However, alveolar epithelial cells contained positive hybridization signals cells at 1,3 and 7 PID. Cells containing viral RNA were detected from 1 to 11 PID, in exudate within bronchial and bronchiolar lumina; and from 3 to 7 PID in

alveolar exudates. Positive hybridization signals were identified in interstitial mononuclear cells and in bronchi associated lymphoid tissue from 3 to 11 PID. The highest signal intensity of positive cells were observed at 3 and 7 PID, coinciding with high virus antibodies titres and with the most important histopathological findings.

ACKNOWLEDGEMENTS

A.J. Masot received a fellowship from the Dirección General de Investigación Científica y Técnica (Ministerio de Educación y Ciencia) of Spain.

This work was supported by the Education, Science and Technology Council of Junta de Extremadura; Project N°: IPR00A005 and by Hoeschst Roussel Agri-Vet Company, USDA Animal Health Funds, the Nebraska Agricultural Experiment Station.

REFERENCES

- AL-DARRAJI, A. M., R. C., CUTLIP, H. D. LEHMKUHL, D. L. GRAHAM. 1982. Experimental infection of lambs with bovine respiratory syncytial virus and *Pasteurella haemolytica*: Pathologic studies. *A. J. Vet. Res.* 42: 224-229.
- BELKNAP, E. B., D. K. CISZEWSKI, J. C. BAKER. 1995. Experimental respiratory syncytial virus infection in calves and lambs. *J. Vet. Diagn. Invest.* 7: 285-298.
- BRUHN, F. W., A. S. YEAGER. 1977. Respiratory syncytial virus in early infancy. Circulating antibody and the severity of infection. *A. J. Dis. Child.* 131: 145-148.
- BRYSON, D. G., P. F. CUSH, M. S. MCNULTY, M. PLATEN, G. M. ALLAN. 1988. An immunoperoxidase method of detecting respiratory syncytial virus antigens in paraffin sections of pneumonic bovine lung. *A. J. Vet. Res.* 49: 1121-1126.
- BRYSON, D. G., S. MCCONELL, M. MCALISKE, M. S. MCNULTY. 1991a. Ultrastructural features of alveolar lesions in induced respiratory syncytial virus pneumonia of calves. *Vet. Pathol.* 28: 286-292.
- BRYSON, D. G., M. F. PLATTEN, S. MCCONELL, M. S. MCNULTY. 1991b. Ultrastructural features

- of lesions in bronchiolar epithelium in induced respiratory syncytial virus pneumonia of calves. *Vet. Pathol.* 28: 293-299.
- CASTLEMAN, W. L., J. C. LAY, E. J. DUBOVI, D. O. SLAUSON. 1985. Experimental bovine respiratory syncytial virus infection in conventional calves: light microscopic lesions, microbiology and studies on lavaged lung cells. *A. J. Vet. Res.* 46: 547-553.
- CIRINO, N. M., J. R. PANUSKA, A. VILLANI, H. TARAF, N. REBERT, R. MEROLLA, P. TSIVITSE, I. A. GILBERT. 1993. Restricted replication of respiratory syncytial virus in human alveolar macrophages. *J. Gen. Virol.* 74: 1527-1537.
- CISZEWSKI, D. K., J. C. BAKER, R. F. SLOCOMBE, J. F. REINDEL, D. M. HAINES, E. G. CLARK. 1991. Experimental reproduction of respiratory tract disease with bovine respiratory syncytial virus. *Vet. Microbiol.* 28: 39-60.
- COWAN, S.T. 1974. Cowan and Steel's. Manual for Identification of Medical Bacteria, Cambridge, UK: University Press, Cambridge.
- ELAZHARY, M., A. SILIM, R. S. ROY. 1981. Interferon, fluorescent antibody, and neutralizing antibody responses in sera of calves inoculated with bovine respiratory syncytial virus. *A. J. Vet. Res.* 42: 1378-1382.
- KORBECKI, M., M. MAKSYMOWICZ. 1977. An attempt to determine the relation between levels of IgG, IgA and IgM immunoglobulins and titres of antibodies against RS virus. *Arch. Immunol. Ther. Exp.* 25: 477-482.
- LEHMKUHL, H. D., P. M. GOUGH, D. E. REED. 1979. Characterization and identification of a bovine respiratory syncytial virus isolated from young calves. *A. J. Vet. Res.* 40: 124-126.
- LERCH, R.A., K. ANDERSON, V. L. AMANN, G. W. WERTZ. 1991. Nucleotide sequence analysis of the bovine respiratory syncytial virus fusion protein mRNA and expression from recombinant vaccinia virus. *Virology.* 181: 118-131 .
- MASOT, A. J., L. GÓMEZ, C. GÓMEZ-TEJEDOR, S. MARTÍNEZ, E. REDONDO. 1993a. The immunologic response to experimental infection with bovine respiratory syncytial virus in lambs. *J. Vet. Med. B.* 40: 659-666.
- MASOT, A. J., C. GÓMEZ-TEJEDOR, I. TOVAR, A. GÁZQUEZ, E. REDONDO. 1993b. Location of bovine respiratory syncytial virus antigens in the lung of experimentally-infected lambs: comparative study using indirect fluorescent antibody test, avidin-biotin-peroxidase complex and transmission electron microscopy. *E. J. Histochem.* 37: 75-82.
- MASOT, A. J., A. GÁZQUEZ, S. REGODÓN, A. FRANCO, E. REDONDO. 1995. Lesions in lambs experimentally infected with bovine respiratory syncytial virus. *Histol. Histopathol.* 10: 71-77.
- MASOT, A. J., C. GÓMEZ-TEJEDOR, L. GÓMEZ, A. GÁZQUEZ, E. REDONDO. 1996. Pathological study of experimentally induced bovine respiratory syncytial virus in lambs. *J. Vet. Med. B.* 43: 233-243.
- MATHEISE, J. P., K. WALRAVENS, A. COLLARD, P. COPPE, J. J. LETESSON. 1995. Antigenic analysis of the F protein of the bovine respiratory syncytial virus: identification of two distinct antigenic sites involved in fusion inhibition. *Archiv. Virol.* 140: 993-1005.
- MOHANTY, S. B., A. L. INGLING, M. G. LILLIE. 1975. Experimentally induced respiratory syncytial viral infection in calves. *A. J. Vet. Res.* 36: 417-419.
- OBERST, R. D., M. P. HAYS, K. J. HENNESSY, L. C. STINE, J. F. EVERMANN, C. KELLING. 1993. Identifying bovine respiratory syncytial virus by reverse transcription-polymerase chain reaction and oligonucleotide hybridizations. *J. Clin. Microbiol.* 31: 1237-1240.
- PASTEY, M. K., S. K. SAMAL. 1993. Structure and sequence comparison of the respiratory syncytial virus fusion protein. *Virus Res.* 29: 195-202.
- PATON, A. W., J. C. PATON, A. J. LAWRENCE, P. N. GOLDWATER, R. J. HARRIS. 1992: Rapid detection of respiratory syncytial virus in nasopharyngeal aspirates by reverse transcription and polymerase chain reaction amplification. *J. Clin. Microbiol.* 30: 901-904.
- PIRIE, H. M., L. PETRIE, C. R. PRINGLE, E. M. ALLAN, J. G. KENNEDY. 1981. Acute fatal pneumonia in calves due to respiratory syncytial virus. *Vet. Rec.* 108: 411-416.
- REDONDO, E., A. J. MASOT, S. MARTÍNEZ, A. JIMÉNEZ, A. GÁZQUEZ. 1994: Spontaneous bovine respiratory syncytial virus infection in goats: pathological findings. *J. Vet. Med. B* 41: 27-34.
- SCHRIJVER, R. M., J. A. KRAMPERS, W. G. J. MIDDEL, J. P. M. LANGEDIJK, J. T. VAN OIRSCHOT. 1994: Bovine respiratory syncytial virus replicates minimally in bovine alveolar macrophages. *Archiv Virol.* 140: 1905-1917.
- SUR, J. H., V. L. COOPER, J. A. GALEOTA , R. A. HESSE, A. R. DOSTER, F. A. OSORIO. 1996. *In vivo* detection of porcine reproductive and respiratory syndrome virus RNA in situ hybridization at different times postinfection. *J. Clin. Microbiol.* 34: 2280-2286.

- THOMAS, L. H., E. J. STOTT, A. P. COLLINS, S. CROUCH, J. JEBBETT. 1984. Infection of gnotobiotic calves with bovine and human isolate of respiratory syncytial virus. Modification of the response by dexamethasone. *Arch. Virol.* 79: 67-77.
- VIUFF, B., A. UTTENTHAL, C. TEGTMEIER, S. ALEXANDERSEN. 1996. Sites of replication of bovine respiratory syncytial virus in naturally infected calves as determined by *in situ* hybridization. *Vet. Pathol.* 33: 383-390.
- WALRAVENS, K., R. KETTMANN, A. COLLARD, P. COPPE, A. BURNY. 1990. Sequence comparison between the fusion protein of human and bovine respiratory syncytial viruses. *J. Gen. Virol.* 71: 3009-3041.
- WALSH, E. E., M. V. BRANDRISS, J. J. SCHLESINGER. 1985. Purification and characterization of the respiratory syncytial virus fusion protein. *J. Gen. Virol.* 66: 409-415.

