

## Effects of plant extracts on the growth of beneficial indigenous lactic acid bacteria (BLAB) for their potential use in preventing bovine reproductive tract infections

María Hortencia Miranda<sup>a</sup>, María Elena Fátima Nader-Macías<sup>a\*</sup>

**ABSTRACT.** There is a renewed interest in products based on phytochemicals, prebiotics and probiotics applied to different hosts to exert effects of immunomodulation, anti-inflammation and analgesia. The microbiome of the bovine reproductive tract can become unbalanced for many reasons, favoring the entry and proliferation of pathogenic microorganisms, currently treated with antibiotics that exert adverse effects and generate antimicrobial resistance. To deal with this situation, “phytobiotic” formulas are proposed that combine phytochemicals and probiotics. This work aims to study the effect of plant extracts, prebiotics and vitamins on the growth of native beneficial lactic acid bacteria (BLAB), to be further potentially applied in the design of phytobiotic formulas. Nine beneficial strains isolated from different bovine ecosystems were evaluated against nine phytochemicals, two prebiotics and five vitamins. Affinity was assayed using the diffusion technique on agar plates, and the effect of the phytochemicals on the growth of lactic acid bacteria by microplates. The growth of all the strains was affected by some plant extracts, showing a stimulating or inhibitory effect. Diffusion-agar plates show that only vitamin A affected the viability of *Lactobacillus johnsonii* CRL1702 at concentrations higher than 7.5 mg/ml. When studying the growth kinetics of the strains with the phytochemicals, the results show that the effect was different in each of one the associated strains + plant extracts, indicating a strain-specific effect of plant extracts on each BLAB strain. Lapacho and Malva stimulate the growth of most microorganisms, then were selected to be combined with BLAB to design a phytobiotic formula with potential therapeutic activity to treat bovine reproductive infections. Plant extracts at the evaluated concentrations did not inhibit the growth of most of the pathogens responsible for endometritis. On the other hand, the highest concentrations of phenolic compounds were detected in Echinacea, Lapacho and Llantén; and the best percentages of antioxidant activity were evidenced in Garlic, Blueberry and Chamomile (<60%).

**Keywords:** Probiotic lactic bacteria, *Handroanthus impetiginosus*, *Malva sylvestris*, phytobiotics, bovine reproductive health.

### INTRODUCTION

The autochthonous microbiota of the different bovine tracts and mucosa are constituted by a wide variety of microorganisms in ecological balance, called “microbiome”, which is currently considered one more organ since it provides metabolic activities, coding capacities and fulfills a fundamental role and variety of physiological functions including immunomodulation and prevention of infection in humans and animals (Li *et al.* 2013; Stumpf *et al.* 2013). Lactic acid bacteria (LAB) are one of the most significantly important groups of bacteria in the food industry. They have long been consumed in dairy products all around the world and most of them are classified as “Generally Recognized As Safe” (GRAS) and into Qualified Presumption of Safety (QPS) microorganisms, because they are non-pathogenic, suitable for technological and industrial processes, having the ability to produce many antimicrobial substances (Reuben *et al.*, 2020; Shehata *et al.*, 2016). In the last decade, LAB has received increased attention and are

widely used as probiotics, defined as “live microorganisms that exert health benefits on the host when ingested in adequate amounts” (FAO/WHO, 2006). Furthermore, lactobacilli are also under development as delivery systems for vaccines and treatments (Sun *et al.*, 2015).

On the other hand, there is a growing interest in natural ingredients, including plant sources and derivatives, both from consumers and producers in the food and pharmaceutical industries. People search into the market for products that are artificial and synthetic additives-free to promote their health and animal sanity (Marchesi *et al.*, 2020). In almost all countries, plants have been widely used throughout history for the treatment and prevention of different diseases and infections in humans and domestic animals. Today, these traditional treatments are recommended in human and veterinary medicine due to their promising therapeutic efficacy with minimal side effects and reduction of chemotherapeutic drug residues in animal products consumed by humans (Gurib-Fakim, 2006). Most of them are included in the Pharmacopoeia regulations, while the perspective and future approaches of ethnopharmacological research are developed in parallel with advances in clinical and laboratory sciences, mainly phytochemistry and pharmacology (González-Stuart *et al.*, 2017). Medicinal plants with their well-established history are an excellent resource of natural products used as an alternative therapy (Mushtaq *et al.*, 2018). The limitation of therapeutic options for emerging multi-resistant microorganisms and the urgent need for new (or old uses)

Received: 11.10.2022

Accepted: 20.03.2023

<sup>a</sup>Laboratorio Farmabióticos, Centro de Referencia para Lactobacilos, Consejo Nacional de Investigaciones Científicas y Técnicas. San Miguel de Tucumán, Tucumán, Argentina.

\*Corresponding author: fnader@cerela.org.ar, fatynader@gmail.com

natural and safe combinations are emerging. Then, the concept of phytobiotics was conceived, to combine safe phytoderivatives and probiotic bacteria for the design of new formulations, to advance in the combination/synergy of the immunomodulatory, anti-inflammatory, anticancer, analgesic and antioxidant effects reported individually in both, LAB probiotics and plant extracts (Nader Macías *et al.*, 2008; Ayrle *et al.*, 2016; Martínez & Lujan, 2011). These bioactive ingredients must be included in formulas in such a way that they must be compatible, protect against harsh environmental and process conditions, and be safely delivered to cells and target organs. In the area of probiotics, our research group has a long history of experiments carried out *in vitro* or in animal models, complemented by studies on the design of different types of vaginal probiotic formulas (Ocaña *et al.*, 1999; De Gregorio *et al.*, 2016; Leccese Terraf *et al.*, 2014).

Therefore, the objective of this work was to evaluate the effect of phytoextracts (powdered) on the growth of bovine homologous vaginal probiotic lactobacilli and their affinity/compatibility to select those to be combined for the further design of a phytobiotic formulas. Also, to determine the phytocompounds effect on bovine vaginal pathogens, and the phenolic content and antioxidant activity of the plant extracts, to define if are related with the beneficial effect. The best combinations were defined to advance in the design of an intravaginal formula to be further applied to the reproductive health of postpartum cows.

## MATERIALS AND METHODS

### BENEFICIAL AUTOCHTHONOUS LACTIC ACID BACTERIA (BLAB) AND GROWTH CONDITIONS

The bacterial strains used in this study were beneficial lactobacilli previously isolated from three bovine ecosystems: the vaginal tract and mammary gland of cows and the gastrointestinal tract of calves (Tucumán, Argentina). The strains used are included in Table 1, together with the previously determined beneficial properties (Otero *et al.*, 2006; Espeche *et al.*, 2012; Maldonado *et al.*, 2012; Miranda *et al.*, 2020). The microorganisms were preserved in yeast extract milk (13% skim milk, 0.5% yeast extract and 1% glucose) at -20 °C, subcultured 3 times in MRS broth (De Man, Rogosa and Sharpe, Biokar Diagnostics, Beauvais, France) and incubated for 12 hours at 37 °C. Protocols were performed with the third subculture at 2%. The vaginal tract strains were isolated from vaginal brushing samples (Otero *et al.*, 2006), from the mammary gland/milk (Espeche *et al.*, 2012) adult cows. Calves LAB were obtained from saliva and gastrointestinal tract (faeces) (Maldonado *et al.*, 2012).

### PREPARATION OF PLANT EXTRACTS, VITAMINS AND PREBIOTICS

The solid vegetable compounds evaluated were obtained from SAPORITI (Buenos Aires, Argentina) and selected

**Table 1.** Beneficial characteristics of the lactic acid bacteria strains selected and evaluated.

Microorganism	Beneficial properties*	Isolation Origin
<i>Lactobacillus gasseri</i> CRL 1412	H <sub>2</sub> O <sub>2</sub> producer, hydrophobic, biofilm former and EPS producer	Bovine vagine*
<i>Lactobacillus gasseri</i> CRL 1421	H <sub>2</sub> O <sub>2</sub> and lactic acid producer, hydrophobic, biofilm-forming	
<i>Lactobacillus gasseri</i> CRL 1460	Hydrophobic	
<i>Lactobacillus gasseri</i> CRL 1461	Lactic acid producer. Hydrophobic, self-aggregating, high adhesion capacity	
<i>Lactococcus lactis</i> subsp. <i>lactis</i> CRL 1656	H <sub>2</sub> O <sub>2</sub> and lactic acid producer. Inhibits <i>S. aureus</i> , <i>S. uberis</i> , <i>S. agalactiae</i> and SCN	Bovine mammary gland <sup>γ</sup>
<i>Pediococcus pentosaceus</i> CRL 1831	H <sub>2</sub> O <sub>2</sub> producer, biofilm former, EPS +	
<i>Weisella cibaria</i> CRL 1833	Hydrophobic, biofilm-forming, EPS +	
<i>Lactobacillus johnsonii</i> CRL 1693	Lactic acid producer. Self-aggregating, hydrophobic, inhibits <i>E. coli</i> , <i>S. aureus</i> , <i>S. dublin</i> .	Faecal matter calves <sup>+</sup>
<i>Ligilactobacillus murinus</i> CRL1695	H <sub>2</sub> O <sub>2</sub> and lactic acid producer. Self-aggregating, biofilm-forming, inhibits <i>S. typhimurium</i>	
<i>Limosilactobacillus mucosae</i> CRL1696	H <sub>2</sub> O <sub>2</sub> producer	
<i>Ligilactobacillus salivarius</i> CRL1702	Lactic acid producer. Hydrophobic, inhibits <i>S. typhimurium</i>	

CRL: Centro de Referencia para Lactobacillus

\*Published previously by Otero *et al.* (2006).

<sup>γ</sup>Published previously by Espeche *et al.* (2012).

<sup>+</sup>Published previously by Maldonado *et al.* (2012).

<sup>\*</sup>Published previously by Miranda *et al.* (2020).

from the list of natural extracts approved by different Pharmacopoeias (Argentinian Pharmacopoeia; European Pharmacopoeia) for human and veterinary applications, included in Table 2 with their ethnopharmacological properties: *Allium sativa* (Garlic), *Vaccinium myrtillus* (Blueberry), *Atropa belladonna* (Belladonna), *Echinacea angustifolia* (Echinaceae), *Matricaria recutita* (Chamomile), *Handroanthus impetiginosus* (Lapacho), *Malva sylvestris* (Malva), *Larrea divaricata* (Jarilla) and *Plantago major* (Llantén).

#### COMPATIBILITY OF BLAB WITH PLANT EXTRACTS BY THE AGAR PLATE DIFFUSION TECHNIQUE

The Minimal Inhibitory Concentration (MIC) of the plant extract was determined by the agar plate diffusion technique, by using a range between 1.8 and 30 mg/ml.

They were prepared at 60 mg for the microplate assay, dissolved in 1 ml of 25% alcohol, and stored at 4 °C protected from light until use. The vitamins used were: A, B5, B7 and B12 (ICN Biomedicals Inc, Ohio, USA), while powdered corn syrup (Arcor, Lules, Tucuman, Argentina) and inulin (Sigma-Aldrich, USA) was used as prebiotic and dissolved in water (30 mg/ml).

MRS (De Man, Rogosa and Sharpe) agar plates (Biokar Diagnostics, Beauvais, France) (1% agar) were prepared by inoculating 100 µl of different BLAB ( $1 \times 10^6$  and  $1 \times 10^8$  CFU/ml) added to the melted and cooled agar. Once solidified, 25 µl (per spot) of the seriated dilutions of plant extracts, vitamins and prebiotics were inoculated, maintained quiescent until complete diffusion on the agar, and incubated at 37 °C for 48 hours. The inhibition zones (in millimeters) were determined by using a manual caliper after. Assays were performed in triplicate.

**Table 2.** Application/Effect of vegetal extracts (approved in pharmacopoeias) assayed

Scientific name	Vulgar name	Application*	References	Pharmacopoeia
<i>Allium sativum</i>	Garlic	Modulation of the immune system and inflammation. GI tract: antidiarrheal Antibacterial - Anti-inflammatory Insect Bites - Injuries and Wounds	Ayrle <i>et al.</i> , 2016; Mirabeau 2012; Harris, 2001; Pittler, 2007; Iciek, 2009; Oosthuizen, 2017; Anwar 2017; Cheng, 2018	Argentina 8th edition
<i>Malva sylvestris</i>	Malva	Modulation of the immune system and inflammation - Intestinal colic - Wounds and injuries. Mastitis - Intestinal colic - Eye lesions - Antioxidant	Ayrle <i>et al.</i> , 2016; Martínez & Lujan, 2011; Bensaad, 2016; Prudente, 2017; Martins, 2017; Lasparetto, 2011; Vahabi, 2019	Argentina 1er annex
<i>Larrea divaricata</i>	Jarilla	Bumps or swelling - Muscle pain - Placental retention	Martínez & Lujan, 2011; Canale, 2018; Davicino, 2007; Palacio, 2008, 2012; Davicino, 2008; Martino, 2010, 2011; Pedernera, 2006	Argentina 8th edition
<i>Plantago major</i>	Llantén	Shock/inflammation - Wounds and injuries - Antiulcer - Anti-inflammatory - Antidiarrheal - Antitumor.	Martínez & Lujan, 2011; Amos, 2017; Najafian, 2018; Chiang, 2003a, b; Huang, 2009	Argentina 8th edition
<i>Matricaria recutita</i>	Chamomile	Wounds and injuries - Treatment of castration - Antioxidant - Anticancer - Placental retention - Insect and animal bites	Martínez & Lujan, 2011; Miraj, 2016; McKay, 2006; Sharifi-rad, 2018; Park, 2017; Al-Dabbagh, 2019; Mohsenzadeh, 2011	Argentina 8th edition
<i>Echinacea angustifolia</i>	Echinacea	Immune system strengthens and enhancer	Martínez & Lujan, 2011; Barnes, 2005; Arland, 2016; Sharifi-rad, 2018; Sharma, 2009; Stevenson, 2005	European annex 2014
<i>Vaccinium myrtillus</i>	Blueberry	Antioxidants - Anti-inflammatory - Anti-hypertensive - Antimicrobial - Anticancer	Jakesevic, 2013; Korus, 2014; Mykkanen, 2014; Nohynek, 2006; Veljković, 2016	European 8th edition / Argentine Food Code
<i>Atropa belladonna</i>	Belladonna	Antipyretic - Modulates inflammation	Ahmad, 2018; Pedalino, 2004	Argentina 8th edition
<i>Handroanthus impetiginosus</i>	Lapacho	Anti hyper uricemic - Anti-inflammatory - Antioxidant - Anti-tumor - Antiulcerogenic - Analgesic - Anti-allergy	Ferraz-Filha, 2016; Grazziotin, 1992; Kim, 2012; Suo, 2012	Argentina 3rd annex

\*Ethnopharmacological effect of plant extracts<sup>Ω</sup>

## GROWTH KINETICS OF BLAB WITH PHYTOCOMPOUNDS BY MICROPLATE TECHNIQUE

The behavior of 9 plant extracts (SAPORITI drugstore, Buenos Aires-Argentina) at a concentration of 60 mg/ml on 10 different BLAB strains was evaluated: *Lactobacillus gasseri* CRL1412, *Lactobacillus gasseri* CRL1421, *Lactobacillus gasseri* CRL1460, *Lactobacillus gasseri* CRL1461, *Lactococcus lactis* subsp. *lactis* CRL1656, *Pediococcus pentosaceus* CRL1831, *Weissella cibaria* CRL1833, *Lactobacillus johnsonii* CRL1693, *Ligilactobacillus murinus* CRL1695, *Limosilactobacillus mucosae* CRL1696 and *Ligilactobacillus salivarius* CRL1702. The solid plant extracts were weighed and resuspended in 1 ml of 25% ethanol and filtered through sterile 0.22µm membrane filters (Biofil-Syringe Filter). The BLAB inoculum was prepared from the third MRS broth subculture with O.D.<sub>560nm</sub> 0.9-1.0 ( $10^7$ - $10^8$  CFU / ml).

The effect of the plant extracts on the growth of the BLAB strains was evaluated in polystyrene microplates (Extragen-96-well ELISA plate), to which 150 µl of MRS broth (De Man, Rogosa and Sharpe) (Biokar Diagnostics, Beauvais, France) inoculated with 2% BLAB was added. Then 50 µl of the plant extracts were added, gently mixed and incubated at 37 °C for 48 hours. Bacterial growth was determined by optical density (O.D.<sub>560nm</sub>) (Spectronic 20, Bausch and Lomb, Rochester, NY) at 4, 6, 22, 24, 30 and 48 hours. Solvents, individual plant extracts and bacteria in MRS were included as a control. The values of O.D. of bacterial growth were obtained subtracting the O.D. value obtained for each plant extract. All experiments were performed in triplicate.

## EFFECT OF PLANT EXTRACTS ON THE GROWTH OF BOVINE REPRODUCTIVE TRACT PATHOGENS

The effect of 9 plant derivatives against different pathogens isolated from the bovine reproductive tract was evaluated by using the modified agar plate diffusion technique (De Gregorio *et al.*, 2019). *Escherichia coli* 99/14, *Pseudomonas aeruginosa*, *Streptococcus bovis*, *Enterococcus faecalis* and *Listeria monocytogenes* (Nader *et al.*, 2008) were cultured in LAPTg broth (Biokar Diagnostics, Beauvais, France) for 12 hours at 37 °C under microaerophilic conditions, in an anaerobic jar (AnaeroGen Oxoid). Briefly, LAPTg agar plates (15 ml of LAPTg with 1% agar) were inoculated with  $1 \times 10^6$  and  $1 \times 10^8$  CFU/ml of each pathogen (Pasteris *et al.*, 2011). Aliquots (25 µl) of the different solid plant extracts dissolved in 25% hydroalcoholic solution (30, 15, 7.5 and 3.25 mg/ml) were added to the holes (4mm in diameter) aseptically made in the agar plates (in triplicate), incubated later for 24 hours at 37°C. The inhibition halos were quantified in mm, determining the Minimum Inhibitory Concentration (MIC) of each of the plant extracts. The extracts were added also as spots on the surface of the pathogens-agar plates (De Gregorio *et al.*, 2019).

## DETERMINATION OF TOTAL PHENOLIC COMPOUNDS OF PLANT EXTRACTS

The total phenolic content of the phytochemicals was determined by the Zielinski & Kozłowska (2000) technique, with the following modifications: 100 µl of each plant extract (1mg/ml) and 900 µl bidistilled water were mixed with 100 µl Folin-Ciocalteu reagent (Sigma-Aldrich, Darmstadt, Germany) (diluted to 50% in bidistilled water). After 2 minutes at room temperature, 400µl of a Na<sub>2</sub>CO<sub>3</sub> solution (Ciccarelli, Santa Fe, Argentina) (15.9%, in bidistilled water) was added and left to stand for 10 minutes in the dark. Then 200 µl were seeded in microplates (in triplicate). Absorbance was measured in a microplate reader at 765 nm in a UV-VIS spectrophotometer (Spectronic 20, Bausch and Lomb, Rochester, NY). The calibration curve was prepared with gallic acid (GA) (Sigma-Aldrich; Merck KGaA, Darmstadt, Germany) in ethanol as a standard, in a concentration range of 2.5 to 1000 µg. The regression equation used was  $y = 0.0395x + 0.1633$ , and the correlation coefficient  $R^2 = 0.9907$ . The results were expressed in mg of GAE extract/ml.

## ANTIOXIDANT ACTIVITY OF PLANT EXTRACTS

The antiradical activity was evaluated by the photometric method, according to Wu *et al.* (2003). The DPPHu radical (1,1-diphenyl-2-picryl hydrazyl, Sigma-Aldrich, Darmstadt, Germany) was dissolved in 80% methanol up to an absorbance equal to 1.0 at 517 nm. 100 µl aliquots of each phytochemicals (1mg/ml) were mixed with 200 µl of DPPHu. The decrease in absorbance was determined in a microplate spectrophotometer (Spectronic 20, Bausch and Lomb, Rochester, NY) at 517 nm, with readings every 30 seconds for 10 minutes. The percentage of inhibition of DPPHu radical was calculated through the equation described by Burda & Oleszek (2001).

$$AAR = 100 \times (1 - A_m/A_0)$$

Where  $A_m$  is the absorbance of the mixture at an infinite time and  $A_0$  is the absorbance of the DPPH solution before the addition of the samples.


## STATISTICAL ANALYSIS

The results were subjected to analysis of variance (ANOVA) and a significance level of 5% ( $p \leq 0.05$ ) was applied, using the Minitab analytical program.

## RESULTS

## BLAB COMPATIBILITY WITH PLANT EXTRACTS

The diffusion method on MRS agar plates determined that all plant extracts assayed are compatible with BLAB, as



**Figure 1.** Inhibition of vitamin A on *Ligilactobacillus salivarius* CRL1702:  $10^6$  CFU/ml (a) and  $10^8$  CFU/ml (b). The black circles indicate the inhibition spots.

well as inulin and corn syrup used as prebiotics, except *L. salivarius* CRL1702 ( $10^6$  and  $10^8$  CFU/ml) (Figure 1A-B) which was inhibited by vitamin A at 30 and 15 mg/mL concentrations (MIC = 7.5 mg/mL).

#### BLAB GROWTH IN THE PRESENCE OF DIFFERENT PLANT EXTRACTS

The plant extracts under study affected the growth of many BLAB strains since they were either stimulated or inhibited. Figure 2 shows the effect of different phytochemicals on BLAB growth, indicating that the effect is strain-specific. The claimed pharmacological effects of all the phytoderivatives assayed (Table 2), were not related to the degree of inhibition exerted on the growth of the strains; the growth of each BLAB with the added phytochemicals, evaluated by O.D.<sub>560nm</sub>, indicated that the effect is different in each one of the strains + plant extracts associations. Some examples are shown in Figure 2: a-g: represents obligate heterofermentative bacteria (OHe) and, h-i: obligate homofermentative lactobacilli (OHO). The growth in standard medium (MRS) of each strain is represented by a dotted line. Figure 2a shows the behavior of *L. gasseri* CRL1412 stimulated by Lapacho, Malva, Llantén and Manzanilla during the first 6 hours, but inhibited in a higher degree by Blueberry, Belladonna, Echinacea and Garlic. However, it is strongly stimulated by Jarilla starting at 12 hours. *L. gasseri* CRL1421 follows a growth pattern similar to CRL1412, showing good growth together with Lapacho and Malva, but inhibited by Belladonna, Llantén and Echinacea (Figure 2b). Figure 2c indicated the growth kinetics of *L. gasseri* CRL1460 strongly stimulated by Lapacho and Malva at 3 and 6 hours, respectively. After 6 hours, this strain was inhibited by most of the plant extracts, except Jarilla. *L. gasseri* CRL1461 (Figure 2d) which showed the highest growth between 3 and 6 hours. Malva, Manzanilla, Lapacho, Llantén and Equinacea produced even higher growth than in the standard broth


(MRS broth). Lapacho, Malva and Llantén facilitated the growth of *L. lactis* subsp. *lactis* CRL1656 during the first 6 hours. Malva reaches a maximum O.D.<sub>540nm</sub> at 24 hours ( $1.43 \pm 0.039$ ), followed by Echinacea with an O.D.<sub>540nm</sub> of  $1.33 \pm 0.10$ ; while Belladonna suppressed the growth (Figure 2e). Figure 2f shows the growth of *P. pentosaceus* CRL1831 with plant extracts, indicating that Garlic and Malva inhibited this strain, while Blueberry, Lapacho and Manzanilla favor its growth at 24 hours. *L. salivarius* CRL1702 shows similar kinetics but only Llantén allowed higher growth rates at 24 h. Blueberry negatively affects both strains (Figure 2g).

The growth of *W. cibaria* CRL1833 is represented in Figure 2h, where Blueberry, Garlic, Jarilla, Belladonna and Echinacea were showed to inhibit it, while Lapacho and Malva strongly stimulated. *L. mucosae* CRL1696 evidenced an inhibitory pattern of all the plant extracts during the first 6 hours, however, Malva, Llantén, Lapacho and Echinacea produced a higher growth than in MRS at 24 hours (Figure 2i).

The degree of interaction between BLAB and plant extracts is summarized in Figure 4, showing the compatible or non-compatible combinations: stimulating (green) or inhibitory (red) effect of the phytochemicals on the growth of each strain. Gray boxes indicate “No effect” because there is no modification of the growth with the phytochemicals when compared to the MRS control.

#### EFFECT OF PLANT EXTRACTS ON THE GROWTH OF UROGENITAL PATHOGENS

The plant extracts assayed at different concentrations was not able to inhibit the growth of most of the frequent pathogenic microorganisms that cause bovine metritis, although the antimicrobial effect of many of them was previously reported (Oosthuizen *et al.*, 2017; Martínez & Lujan 2011; Nohynek *et al.*, 2006; Sharifi-Rad *et al.*, 2018; Canale *et al.*, 2018).



**Figure 2.** Effect of plant extracts on the growth of obligate heterofermentative (a-g) and obligate homofermentative (h-i) BALB. The asterisk (\*) indicates significant differences ( $p < 0.05$ ) in the growth of the strain in different combinations of MRS + plant extract tested according to Tukey's test.

QUANTIFICATION OF TOTAL PHENOLIC COMPOUNDS (TPC) AND ANTIOXIDANT ACTIVITY IN PLANT EXTRACTS

The results obtained by evaluating the content of TPC and the antioxidant activity of the nine plant extracts were determined to select those with higher antioxidant activity, to promote the beneficial effect when combined with probiotics to the phyto-biotic formula. Echinacea is the plant extract with the highest concentration of phenolic compounds. Lapacho and Jarilla did not show significant differences between them, or between Garlic and Belladonna. On the other hand, Blueberry and Malva did not present significant differences between them, showing the lowest TPC values (Figure 3a). Garlic, Blueberry, and Chamomile recorded the highest percentages of antioxidant activity, without significant differences between them, followed by Malva. Belladonna was the phyto-compound with the lowest antioxidant activity (Figure 3b).

DISCUSSION

There is increasing interest in natural ingredients, including plant sources and their derivatives, and beneficial microorganisms from both consumers and producers in the food and pharmaceutical industries. People search the market for products and foods free of artificial additives that can promote their health, many of them with a very long application based on concepts or uses derived from

custom. The same interest is shown in animal production systems.

On the other hand, the use of lactic acid bacteria as probiotics in different mucosal sites is widely recommended (ISSAP, 2016; McFarland *et al.*, 2018). Specifically, in the bovine reproductive system, different scientists have shown that the application of an adequate combination of probiotics could effectively counteract an endometrial infection by *E. coli* (Genis *et al.*, 2016, 2017) and improve productive parameters (Ametaj *et al.*, 2014). Then, to increase/enhance beneficial effects (such as immunomodulation, and inhibition of pathogenic microorganisms), it is possible to combine them with extracts from plants currently or historically applied to the urogenital tract to design different formulas intended to exert or produce a synergistic effect on the host. Therefore, it is of the utmost importance to determine their compatibility and if the phyto-derivatives affect the growth of BLAB to be used in a single beneficial formula.

In human medicine, some results have been published regarding the administration of a vaginal gel containing *Thymus vulgaris* and *Eugenia caryophyllus* together with two *Lactobacillus* strains specifically formulated in slow-release capsules for the treatment of recurrent bacterial vaginosis or vulvovaginal candidiasis (Murina *et al.*, 2018) however, there are no previous studies that report the compatibility between the two bioactive components of the commercial formula. It is very important, and as a first step for the design of the formula, to evaluate the effect




Figure 3. Quantification of phenolic compounds (a) and anti-oxidant activity (b) in plant extracts.

The different letters represent statistically significant differences (p<0.005) (Tukey’s test).

Vegetable extract \ BALB	Garlic	Blueberry	Belladonna	Echinacea	Chamomile	Hammamelis	Lapacho	Llantén	Malva	Jarilla
<i>L. gasseri</i> CRL1412	Yellow	Red	Orange	Light Green						
<i>L. gasseri</i> CRL1421	Yellow	Light Green	Yellow	Light Green	Light Green	Grey	Light Green	Light Green	Light Green	Light Green
<i>L. gasseri</i> CRL1460	Yellow	Light Green	Orange	Light Green						
<i>L. gasseri</i> CRL1461	Orange	Orange	Yellow	Light Green						
<i>L. lactis</i> subsp. <i>lactis</i> CRL1656	Red	Red	Orange	Light Green						
<i>P. pentosaceus</i> CRL1831	Orange	Light Green	Orange	Light Green						
<i>W. cibaria</i> CRL1833	Orange	Red	Orange	Light Green						
<i>L. mucosae</i> CRL1696	Orange	Orange	Orange	Light Green						
<i>L. salivarius</i> CRL1702	Red	Red	Yellow	Light Green						

Figure 4. Compatibility between beneficial vaginal lactic acid bacteria in the presence of plant extracts. Red: “Inhibitory effect”; Green: “Stimulating effect”; Shades of color indicate greater or lesser inhibition (Red, Orange, Yellow) or stimulation (Dark Green, Light Green) and “No effect” (Grey).

of the phytoderivatives on the growth of lactobacilli. The results of this work show the interactions of a long list of phytoderivatives approved in the Pharmacopoeias to be used for both oral and local administration with bovine BLAB probiotic strains characterized in our laboratory. The compatibility of phytoderivatives and strains of bovine lactic acid bacteria in veterinary medicine has not been previously studied, therefore, the issue is of the utmost importance, and considered as a first step to design a novel formula, to evaluate the effect of phytoderivatives, whether liquid or extract, on the growth of BLAB. The results obtained in this work indicate that the effect of the phytoextracts is dependent on the strain since there are no general rules. Each strain and each extract must be tested to define their optimal or adequate combinations. In this work, the evaluation of their combinations was carried out through MIC tests and growth kinetics, which allowed defining which of them could be adequately combined. Different publications have reported the individual use of lactic acid bacteria and plant extracts as curative and/or preventive treatments in veterinary medicine (Pellegrino *et al.*, 2017; Maldonado *et al.*, 2017; Mansilla *et al.*, 2020; Ayrle *et al.*, 2016; Martínez & Lujan 2011; Lillehoj *et al.*, 2018; Marume *et al.*, 2017). Phytotherapy represents one of the most widely applied non-conventional medicines in both human and veterinary medicine (Hahn *et al.*, 2005). According to Viegi *et al.* (2003), large animals (cattle, horses, sheep, goats and pigs) represent 70.5% of domestic animals treated with herbal remedies, because phytotherapy is mainly used on organic farms to further reduce the use of allopathic medicines. Although the use of phytotherapeutic products is increasing in animals, there are few studies and clinical trials reported in the literature on the therapeutic use of different plants for reproductive disorders in cows (El-Shanawani, 1996; Bruni *et al.*, 1997; Lans *et al.*, 2000; Uncini Manganelli *et al.*, 2001; Alawa *et al.*, 2002; Guarrera, 2005).

As previously indicated, there are no available studies on the combination of phytoderivatives and probiotic lactic acid bacteria and their application in the bovine reproductive system for the prevention of infections, reason by which there is a limitation to discuss/comparing the results obtained in this work with some other scientific reports. For this reason, the beneficial effects of the probiotic microorganisms and the plant extracts assayed in this work are included individually. Supported by the limited information available and the constant search for natural and innocuous products, the combination of probiotics and phytocompounds for veterinary use is an interesting and promising vacancy to study, even though there is a requirement to further assay their combination in the host animal.

The plant extracts evaluated in this work were selected for their variety of pharmacological effects and their use in veterinary medicine. Lapacho (*Handroanthus impetiginosus*) was the plant extract that stimulated the growth of the largest number of BLAB strains evaluated,

with statistically significant differences (at 6 hours). Lapacho is traditionally used in folk medicine and mentioned for its therapeutic effects and included in pharmaceutical forms such as infusions and tinctures, mainly as an analgesic, anti-inflammatory and antitumor agent (Grazziotin *et al.*, 1992). Suo *et al.* (2012) demonstrated in an *in vitro* assay that compounds isolated from Lapacho bark significantly suppressed IL-1 $\beta$  and TNF- $\alpha$  cytokine production without significant cytotoxicity in LPS-stimulated cells, suggesting that Lapacho suppresses production the inflammatory cytokines. Therefore, the combination of Lapacho + BLAB strains is a promising strategy for cattle.

Another phytoderivative, Malva (*Malva sylvestris*), shows a stimulating/promoting effect on *L. gasseri* CRL1412, *L. gasseri* CRL1421, *L. gasseri* CRL1460 and *W. cibaria* CRL1833. Several studies report the use of *Malva sylvestris* for veterinary purposes. The decoctions of whole plants are administered to cattle to treat colic, the leaves applied in compresses have shown high effectiveness in the treatment of mastitis in cattle, infusions and decoctions of aerial flowery parts were used against inflammation, wound infection, diarrhea in calves, its use as a curative for skin, reproductive and nervous disorders was also reported (Gasparetto *et al.*, 2011; Viegi *et al.*, 2003). Martins *et al.* (2017) evaluated the anti-inflammatory activity of the Malva extract and observed an inhibition of the release of proinflammatory mediators. Then the association of Malva with selected BLAB strains results in another interesting combination.

Plantain (*Plantago major*) was able to promote the growth of *L. gasseri* CRL1461, *P. pentosaceus* CRL1831, *L. mucosae* CRL1696 and *L. salivarius* CRL1702. Velasco *et al.* (2006) found that the antibacterial activities of *Plantago major* leaves and seeds were shown to inhibit *Escherichia coli*, *Bacillus subtilis* and *Candida albicans* cultures. The rupture of the cell walls of Gram-positive bacteria and the formation of vesicles in Gram-negative bacteria (Metiner *et al.*, 2012) were observed by electron microscope evaluations.

Jarilla (*Larrea divorcata*) shows a stimulating effect on the growth of vaginal strains, which is interesting, since different publications have shown that extracts of *L. divaricata* have immunomodulatory properties (Davicino *et al.*, 2007, 2010). Jarilla also exhibited *in vitro* antimicrobial effects against pathogens (Quiroga *et al.*, 2004; Stege *et al.*, 2006). Then it is possible to indicate that the behavior of stimulation or inhibition of the extract is directly related and depends on each strain and species of microorganism. The Jarilla is an autochthonous phytoderivative widely used in northern Argentina.

Garlic and Belladonna extracts inhibited the growth, to different degrees, of all strains tested. Although their important therapeutic effects have been studied, it is not possible to combine them with probiotic microorganisms, possibly due to their antimicrobial properties.

Combinations of beneficial lactic acid bacteria and plant extracts for veterinary therapeutic purposes have not been previously reported. Phytobiotic formulations for female vaginal application were studied by combining probiotic lactobacilli with different active ingredients (Marchesi *et al.*, 2020), but no products are marketed in our country. These authors advanced in the selection of viable beneficial lactobacilli for their inclusion in the design of phytobiotic formulas for the health of the female urogenital tract, based on their compatibility with natural compounds with ethnopharmacological properties. The phytocompounds studied in this work did not show to inhibit the growth of pathogenic microorganisms responsible for urogenital infections in cows, even though their antimicrobial effects were reported in the literature. Even though this is not an encouraging result, the definitive effect must be assayed in the host animal, because the *in vitro* experiments, predictive in some aspects, sometimes do not agree with the *in vivo* behavior or clinical/animal studies.

The concentration of phenolic compounds and the antioxidant activity in the extract are not directly correlated with the effect on the growth of BLAB nor with the compatibility between them. Further studies should be carried out to determine if any specific type of component is responsible for the stimulating or inhibitory effect. Lillehoj *et al.* (2018) reported that the beneficial effects of phytoextracts are attributed to their antimicrobial and antioxidant properties, which is not demonstrated in our results. The inclusion of phytoderivatives in the diets of animals should alter and stabilize the intestinal microbiota and thus reduce toxic microbial metabolites in the intestine, due to their direct antimicrobial properties on various pathogenic bacteria, resulting in the alleviation of intestinal challenges and immune stress, which would lead to improved performance (Kim *et al.*, 2015). In ruminants, tannins, included in phenolic compounds, modulate the ruminal and intestinal microbiota and improve the growth of certain bacterial populations. The effects of tannins on the rumen microbiota may be different depending on the molecular nature of these polyphenols (Vasta *et al.*, 2008; Min *et al.*, 2015).

## CONCLUSION

The BLAB strains that showed an affinity with the phytoextracts were: *L. gasseri* CRL1421, *L. gasseri* CRL1460, *L. gasseri* CRL1461 and *W. cibaria* CRL1833 with Lapacho, Malva, Llantén and Jarilla, which can be used to design phytobiotic formulas due to their pharmacological effects (antibacterial, anti-inflammatory, immunomodulatory and healing). These results are original, because the compatibility of natural extracts with probiotic lactic acid bacteria strains was studied, in a way to combine the beneficial effect of the two bioactive compounds in the host, which should be further assayed in *in vivo* protocols.

## COMPETING INTERESTS STATEMENT

The authors declare that they have no competing interests.

## ETHICS STATEMENT

No ethical approval was required in this work, as this is an original article with only bacterial isolates and data. No animal or human samples were employed.

## FUNDING

This work was supported by PICT 2017-4324 and 1187 (FONCYT-ANPCYT-MINCYT) and PIP 744 and 545 from CONICET (Consejo Nacional de Investigaciones Científicas y Técnicas, Argentina).

## REFERENCES

- Alawa, J.P., Jokthan, G.E. & Akut, K. (2002). Ethnoveterinary medical practice for ruminants in the subhumid zone of northern Nigeria. *Preventive Veterinary Medicine*, 54(1), 79-90. [https://doi.org/10.1016/s0167-5877\(01\)00273-2](https://doi.org/10.1016/s0167-5877(01)00273-2)
- Ametaj, B.N., Iqbal, S., Selami, F., Odhiambo, J.F., Wang, Y., Gänzle, M.G., Dunn, S.M. & Zebeli, Q. (2014). Intravaginal administration of lactic acid bacteria modulated the incidence of purulent vaginal discharges, plasma haptoglobin concentrations, and milk production in dairy cows. *Research in Veterinary Science*, 96(2), 365-370. <https://doi.org/10.1016/j.rvsc.2014.02.007>
- Ayrle, H., Mevissen, M., Kaske, M., Nathues, H., Gruetzner, N., Melzig, M. & Walkenhorst, M. (2016). Medicinal plants – prophylactic and therapeutic options for gastrointestinal and respiratory diseases in calves and piglets? A systematic review. *BMC Veterinary Research*, 12(89). <https://doi.org/10.1186/s12917-016-0714-8>
- Bourda, E. & Oleszek, W. (2001). Antioxidant and antiradical activities of flavonoids. *J. Agric. Food Chemistry*, 49(6), 2774-2779. <https://doi.org/10.1021/jf001413m>
- Bruni, A., Ballero, M., & Poli, F. (1997). Quantitative ethnopharmacological study of the Campidano Valley and Urzulei district, Sardinia, Italy. *Journal of Ethnopharmacology*, 57(2), 97-124. [https://doi.org/10.1016/S0378-8741\(97\)00055-X](https://doi.org/10.1016/S0378-8741(97)00055-X)
- Canale, F.P., Davila, S.V., Sasso, C.V., Pellarin, N.W. & Mattar Dominguez, M.A. (2018). Immunization with *Larrea divaricata* Cav. proteins elicits opsonic antibodies against *Pseudomonas aeruginosa* and induces phagocytic activity of murine macrophages. *Microbial Pathogenesis*, 118, 257-267. <https://doi.org/10.1016/j.micpath.2018.03.029>
- Davicino, R., Mattar, A., Casali, Y., Porporatto, C., Correa, S.G. & Micalizzi, B. (2007). *In Vivo* Immunomodulatory Effects of Aqueous Extracts of *Larrea divaricata* Cav. *Immunopharmacology and Immunotoxicology*, 29(3-4), 351-366. <https://doi.org/10.1080/08923970701619703>
- Davicino R., Manuele M.G., Turner S., Ferraro G. & Anesini C. (2010). Antiproliferative activity of *Larrea divaricata* Cav. on lymphoma cell line: participation of hydrogen peroxide in its action. *Cancer Investigation*, 28(1), 13-22. <https://doi.org/10.3109/07357900902849665>
- De Gregorio, P.R., Juárez Tomás, M.S. & Nader-Macías, M.E. (2016). Immunomodulation of *Lactobacillus reuteri* CRL1324 on Group B *Streptococcus Vaginal* Colonization in a Murine. *Experimental Model. American Journal of Reproductive Immunology*, 75(1), 23-35. <https://doi.org/10.1111/aji.12445>
- De Gregorio, P.R., Silva, J.A., Marchesi, A. & Nader-Macías, M.E.F. (2019). Anti-Candida activity of beneficial vaginal lactobacilli in *in vitro* assays and in a murine experimental model. *FEMS Yeast Research*, 19(2). <https://doi.org/10.1093/femsyr/foz008>
- Deng, Q., Odhiambo, J.F., Farooq, U., Lam, T., Dunn, S.M., Gänzle, M.G. & Ametaj, B.N. (2015). Intravaginally administered lactic acid bacteria expedited uterine involution and modulated hormonal

- profiles of transition dairy cows. *Journal of Dairy Science*, 98(9), 6018-6028. <https://doi.org/10.3168/jds.2014-8559>
- El-Shanawani, M.A.A. (1996). *Plants Used in Saudi Traditional Medicine KACST Riyadh*. King Saud University Press, Saudi Arabia.
- Espeche, C., Pellegrino, M., Frola, I., Larriestrac, A. & Bogni, C. (2012). Lactic acid bacteria from raw milk as potentially beneficial strains to prevent bovine mastitis. *Anaerobe*, 18, 103-109. <https://doi.org/10.1016/j.anaerobe.2012.01.002>
- FAO/OMS. (2006). Probiotics in food, healthy and nutritional properties and guidelines for their evaluation. *FAO Food and Nutrition Paper*, (85). WHO/FAO, Rome, Italy.
- Gasparrutto, J.C., Ferreira Martinsa, C.A., Hayashia, S.S., Otuky, M.F. & Pontarolo, R. (2012). Ethnobotanical and scientific aspects of *Malva sylvestris L.*: a millennial herbal medicine. *Journal of Pharmacy and Pharmacology*, 64(2), 172-89. <https://doi.org/10.1111/j.2042-7158.2011.01383.x>
- Genis, S., Bach, A. & Aris, A. (2017). Effects of intravaginal lactic acid bacteria on bovine endometrium: Implications in uterine health. *Veterinary microbiology*. <https://doi.org/10.1016/j.vetmic.2017.04.025>
- Genis, S., Bach, A., Fàbregas, F. & Aris, A. (2016). Potential of lactic acid bacteria at regulating *Escherichia coli* infection and inflammation of bovine endometrium. *Theriogenology*, 85, 625-637. <https://doi.org/10.1016/j.theriogenology.2015.09.054>
- González-Stuart, A. & Rivera, J. (2017). Yellow Oleander Seed, or "Codo de Fraile" (*Thevetia spp.*): A Review of Its Potential Toxicity as a Purported Weight-Loss Supplement. *Journal of Dietary Supplements*, 15(3), 352-364. <https://doi.org/10.1080/19390211.2017.1353565>
- Grazziotin, J.D., Schapoval, E.E.S., Chavesa, C.J., Gleyeb, J. & Henriques, A.T. (1992). Phytochemical and analgesic investigation of *Tabebuia chrysotricha*. *Journal of Ethnopharmacology*, 11, 249-251. [https://doi.org/10.1016/0378-8741\(92\)90051-r](https://doi.org/10.1016/0378-8741(92)90051-r)
- Guarrera, P.M. (2005). Traditional phytotherapy in Central Italy (Marche, Abruzzo, and Latium). *Fitoterapia*, 76(1), 1-25. <https://doi.org/10.1016/j.fitote.2004.09.006>
- Gurib-Fakim, A. (2006). Medicinal plants: traditions of yesterday and drugs of tomorrow. *Molecular Aspects of Medicine*, 27(1), 1-93. <https://doi.org/10.1016/j.mam.2005.07.008>
- Hahn, I., Zitterl-Eglscher, K. & Franz, Ch. (2006). Phytomedicine in dogs and cats: web-based survey among veterinarians in Austria, Germany and Switzerland. *Schweiz Arch Tierheilkd*, 147(3), 135-141. <https://doi.org/10.1024/0036-7281.147.3.135>
- International Scientific Association for Probiotics and Prebiotics (ISAPP). (2016). International Scientific Association for Probiotics and Prebiotics. <https://isappscience.org>.
- Kim, J.E., Lillehoj, H.S., Hong, Y.H., Kim, G.B., Lee, S.H., Lillehoj, E.P. & Bravo, D.M. (2015). Dietary *Capsicum* and *Curcuma longa* oleoresins increase intestinal microbiome and necrotic enteritis in three commercial broiler breeds. *Research in Veterinary Science*, 102, 150-158. <https://doi.org/10.1016/j.rvsc.2015.07.022>
- Lans, C., Harper, T., Georges, K. & Bridgewater, E. (2000). Medicinal plants used for dogs in Trinidad and Tobago. *Preventive Veterinary Medicine*, 45(3-4), 201-220. [https://doi.org/10.1016/S0167-5877\(00\)00123-9](https://doi.org/10.1016/S0167-5877(00)00123-9)
- LecceseTerra, M.C., Mendoza, L.M., Juarez Tomas, M.S., Silva, C. & Nader, M.E.F. (2014). Phenotypic Surface properties (aggregation, adhesion and biofilm formation) and presence of related genes in beneficial vaginal lactobacilli. *Journal of Applied Microbiology*, 117(6), 1761-1772. <https://doi.org/10.1111/jam.12642>
- Lillehoj, H., Liu, Y., Calsamiglia, S., Fernandez-Miyakawa M.E., Chi, F., Cravens, R., Oh, S. & Gay, C. (2018). Phytochemicals as antibiotic alternatives to promote growth and enhance host health. *Veterinary Research*, 49(76). <https://doi.org/10.1186/s13567-018-0562-6>
- Maldonado, N.C., Chiaraviglio, J., Bru, E., De Chazal, L. & Santos, V. (2017). Effect of Milk Fermented with Lactic Acid Bacteria on Diarrheal Incidence, Growth Performance and Microbiological and Blood Profiles of Newborn Dairy Calves. *Probiotics Antimicrob Proteins*, 10(4), 668-676. <https://doi.org/10.1007/s12602-017-9308-4>
- Maldonado, N.C., Silva de Ruiz, C., Otero, M.C., Sesma, F. & Nader-Macías, M.E. (2012). Lactic acid bacteria isolated from young calves--characterization and potential as probiotics. *Research in Veterinary Science*, 92(2), 342-349. <https://doi.org/10.1016/j.rvsc.2011.03.017>
- Mansilla, F., Takagi, M., Garcia-Castillo, V., Aso, H., Nader-Macias, M.E.F., Vignolo, G.M., Kitazawa, H. & Villena, J. (2020). Modulation of Toll-like receptor-mediated innate immunity in bovine intestinal epithelial cells by lactic acid bacteria isolated from feedlot cattle. *Beneficial Microbes*, 11(3), 269-282. <https://doi.org/10.3920/BM2019.0189>
- Marchesi, A., Silva, J., AristimuñoFicoseco, C., Wiese, B. & Nader-Macías, M.E.F. (2020). Effect of phytoderivatives on the growth of homologous beneficial vaginal lactobacilli (BVL) strains and their compatibility for the design of phytobiotics for the vaginal tract health. *Research Square*. <https://doi.org/10.21203/rs.2.22448/v1>
- Martinez, GJ, Lujan, MC (2011). Medicinal plants used for traditional veterinary medicine in the Sierras de Córdoba (Argentina): an ethnobotanical comparison with human medicinal uses. *Journal of Ethnobiology and Ethnomedicine*, 7(1), 23. <https://doi.org/10.1186/1746-4269-7-23>
- Martins, C.A.F., Campos, M.L., Irioda, A.C., Pontarolo Stremel, D., Leal Badaró Trindade, A.C. & Pontarolo, R. (2017). Anti-Inflammatory Effect of *Malva sylvestris*, *Sidacordifolia*, and *Pelargonium graveolens* Is Related to Inhibition of Prostanoid Production. *Molecules*, 22(11), 1883. <https://doi.org/10.3390/molecules22111883>
- Marume, A., Matope, G., Katsande, S., Khoza, S., Mutingwende, I., Mduluz, T., Munodawafa-Taderera, T. & Ndhala, A.R. (2017). Wound Healing Properties of Selected Plants Used in Ethnoveterinary Medicine. *Frontiers in Pharmacology*. <https://doi.org/10.3389/fphar.2017.00544>
- McFarland, L.V., Evans, C.T. & Goldstein, E.J.C. (2018). Strain-Specificity and Disease-Specificity of Probiotic Efficacy: A Systematic Review and Meta-Analysis. *Frontiers in Medicine (Lausanne)*, 5, 124. <https://doi.org/10.3389/fmed.2018.00124>
- Metiner, K., Ozkan, O.O. & Ak, S. (2012). Antibacterial effects of ethanol and acetone Extract of *Plantago major L.* on gram positive & gram negative bacteria. *Kafkas Üniversitesi Veteriner Fakültesi*, 18(3), 503-505. <https://doi.org/10.9775/KVFD.2011.5824>
- Min, Y.W., & Rhee, P-L. (2015). The role of microbiota on the gut immunology. *Clinical Therapeutics*, 37, 968-975. <https://doi.org/10.1016/j.clinthera.2015.03.009>
- Miranda, M.H. & Nader-Macias, M.E.F. (2020). Advances in the Design of a Multi-Strain Homologous Probiotic Formula for Cattle. *Journal of Veterinary Medicine and Animal Sciences*, 3(1), 1022.
- Murina, F., Vicariotto, F. & Di Francesco, S. (2018). *Thymol*, *Eugenol* and lactobacilli in a medical device for the treatment of bacterial vaginosis and vulvovaginal candidiasis. *New Microbiologica*, 41(3), 220-224.
- Mushtaq, S., Shah, A.M., Shah, A., Lone, S.A., Hussain, A., Hassan, Q.P. & Ali, M.N. (2018). Bovine mastitis: An appraisal of its alternative herbal curen. *Microbial Pathogenesis*. <https://doi.org/10.1016/j.micpath.2017.12.024>
- Nader, M.E.F., Otero, M.C., Espeche, M.C. & Maldonado, N.C. (2008). Advances in the desing of probiotic products for the prevention of major diseases in dairy cattle. *Journal of Industrial Microbiology and Biotechnology*. <https://doi.org/10.1007/s10295-008-0438-2>
- Nohynek, L., Alakomi, H.L., Kähkönen, M.P., Heinonen, M., Helander, I.M., Oksman-Caldentey, K.M. & Puupponen-Pimiä, R.H. (2006). Berry Phenolics: Antimicrobial Properties and Mechanisms of Action Against Severe Human Pathogens. *Nutrition and Cancer*, 54, 18-32. [https://doi.org/10.1207/s15327914nc5401\\_4](https://doi.org/10.1207/s15327914nc5401_4)
- Ocaña, V.S., Pesce De Ruiz Holgado, A.A. & Nader-Macías, M.E. (1999). Characterization of a bacteriocin-like substance produced by a vaginal *Lactobacillus salivarius* strain. *Applied and Environmental Microbiology*, 65(12), 5631-5635. <https://doi.org/10.1128%2Faem.65.12.5631-5635.1999>

- Oosthuizen, C., Arbach, M., Meyer, D., Hamilton, C. & Lall, N. (2017). Diallyl Polysulfides from *Allium sativum* as Immunomodulators, Hepatoprotectors, and Antimycobacterial Agents. *Journal of Medicinal Food*. <https://doi.org/10.1089/jmf.2016.0137>
- Otero, M.C., Morelli, L. & Nader Macías, M.F. (2006). Probiotic properties of vaginal lactic acid bacteria to prevent metritis in cattle. *Letters in Applied Microbiology*, 43, 91-97. <https://doi.org/10.1111/j.1472-765X.2006.01914.x>
- Pasteris, S.E., Guidoli, M.G., Otero, M.C., Bühler, M.I. & Nader-Macías, M.E. (2011). *In vitro* inhibition of *Citrobacter freundii*, a red-leg syndrome associated pathogen in raniculture, by indigenous *Lactococcus lactis* CRL 1584. *Veterinary Microbiology*. <https://doi.org/10.1016/j.vetmic.2011.03.025>
- Pellegrino, M., Berardo, N., Giraud, J., Nader-Macías, M.E.F. & Bogni, C. (2017). Bovine mastitis prevention: humoral and cellular response of dairy cows inoculated with lactic acid bacteria at the dry-off period. *Beneficial Microbes*. <https://doi.org/10.3920/BM2016.0194>
- Quiroga, E.N., Sampietro, A.R. & Vattuone, M.A. (2004). *In vitro* fungitoxic activity of *Larrea divaricata* extracts. *Letters in Applied Microbiology*, 39(1), 7-12. <https://doi.org/10.1111/j.1472-765X.2004.01521.x>
- Reuben, R.C., Roy, P.C., Sarkar, S.L., Rubayet, U.I., Alam, A.S.M. & Jahid, I.K. (2020). Characterization and evaluation of lactic acid bacteria from indigenous raw milk for potential probiotic properties. *Journal of Dairy Science*. <https://doi.org/10.3168/jds.2019-17092>
- Sharifi-Rad, M., Mnayer, D. & Morais-Braga, M.F.B. (2018). Echinacea plants as antioxidant and antibacterial agents: From traditional medicine to biotechnological applications. *Phytotherapy Research*. <https://doi.org/10.1002/ptr.6101>
- Shehata, M.G., El Sohaimy, S.A., El-Sahn, M.A. & Youssef, M.M. (2016). Detección de bacterias de ácido láctico probióticas potenciales aisladas para la propiedad de reducción del colesterol y la actividad hidrolasa de sales biliares. *Annals of Agricultural Sciences*, 61, 65-75. <https://doi.org/10.1016/j.aoas.2016.03.001>
- Stege, P.W., Davicino, R.C., Vega, A.E., Casali, Y.A., Correa, S. & Micalizzi, B. (2006). Antimicrobial activity of aqueous extracts of *Larrea divaricata* Cav (jarilla) against *Helicobacter pylori*. *Phytomedicine*, 13(9-10), 724-727. <https://doi.org/10.1016/j.phymed.2005.06.008>
- Sun, Z., Harris, H.M.B., McCann, A., Guo, C., Argimón, S., Zhang, W., Yang, X., Jeffer, I.B., Cooney, J.C., Kagawa, T.F., Liu, W., Song, Y., Salvetti, E., Wrobel, A., Rasinkangas, P., Parkhill, J., Rea, M.C., O'Sullivan, O., Ritari, J., Douillard, F.P., Ross, R.P., Yang, R., Briner, A.E., Felis, G.E., de Vos, W.M., Barrangou, R., Klaenhammer, T.R., Caufield, P.W., Cui, Y., Zhang, H., & O'Toole, P.W. (2015). Expanding the biotechnology potential of lactobacilli through comparative genomics of 213 strains and associated genera. *Nature Communications*, 6, 8322. <https://doi.org/10.1038/ncomms9322>
- Suo, M., Isao, H., Kato, H., Takano, F. & Ohta, T. (2012). Anti-inflammatory constituents from *Tabebuia avellanae*. *Fitoterapia*, 83(8), 1484-1488. <https://doi.org/10.1016/j.fitote.2012.08.014>
- Uncini Manganelli, R.E., Camangi, F. & Tomei, P.E. (2001). Curing animals with plants: traditional usage in Tuscany (Italy). *Journal of Ethnopharmacology*. [https://doi.org/10.1016/S0378-8741\(01\)00341-5](https://doi.org/10.1016/S0378-8741(01)00341-5)
- Vasta, V., Makkar, H.P.S., Mele, M. & Priolo, A. (2008). Ruminal biohydrogenation as affected by tannins *in vitro*. *British Journal of Nutrition*, 102, 82-92.
- Velasco-Lezama, R., Tapia-Aguilar, R., Román-Ramos, R., Vega-Ávila, E. & Pérez-Gutiérrez, M.S. (2006). Effect of *Plantago major* on cell proliferation *in vitro*. *Journal of Ethnopharmacology*, 103(1), 36-42. <https://doi.org/10.1016/j.jep.2005.05.050>
- Viegi, L., Pieroni, A., Guarrera, P.A. & Evangelista, R. (2003). A review of plants used in folk veterinary medicine in Italy as basis for a databank. *Journal of Ethnopharmacology*, 89, 221-244. <https://doi.org/10.1016/j.jep.2003.08.003>
- Wu, H.C., Shiau, C.Y., Chen, H.M. & Chiou, T.K. (2003). Antioxidant activities of carnosine, anserine, some free amino acids and their combination. *Journal of Food and Drug Analysis*, 11(2), 148-153.
- Zieliński, H. & Kozłowska, H., (2000). Antioxidant activity and total phenolics in selected cereal grains and their different morphological fractions. *Journal of Agricultural and Food Chemistry*, 48(6), 2008-2016. <https://doi.org/10.1021/jf990619o>